

LAYERED RESTRICTIONS AND CHEBYSHEV POLYNOMIALS

TOUFIK MANSOUR* AND ALEK VAINSHTEIN†

* Department of Mathematics

† Department of Mathematics and Department of Computer Science
University of Haifa, Haifa, Israel 31905

tmansur@study.haifa.ac.il, alek@mathcs.haifa.ac.il

ABSTRACT. A permutation is called layered if it consists of the disjoint union of substrings (layers) so that the entries decrease within each layer, and increase between the layers. We find the generating function for the number of permutations on n letters avoiding $(1, 2, 3)$ and a layered permutation on k letters. In the most interesting case of two layers, the generating function depends only on k and is expressed via Chebyshev polynomials of the second kind.

2000 MATHEMATICS SUBJECT CLASSIFICATION: Primary 05A05, 05A15; Secondary 30B70, 42C05

1. INTRODUCTION AND MAIN RESULT

Let $\pi \in \mathfrak{S}_n$ and $\tau \in \mathfrak{S}_k$ be two permutations. An *occurrence* of τ in π is a subsequence $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that $(\pi_{i_1}, \dots, \pi_{i_k})$ is order-isomorphic to τ ; in such a context τ is usually called a *pattern*. We say that π *avoids* τ , or is τ -*avoiding*, if there is no occurrence of τ in π . The set of all τ -avoiding permutations in $\mathfrak{S}_n$ is denoted $\mathfrak{S}_n(\tau)$. For an arbitrary finite collection of patterns T , we say that π avoids T if π avoids any $\tau \in T$; the corresponding subset of $\mathfrak{S}_n$ is denoted $\mathfrak{S}_n(T)$.

In several recent papers [CW, MV, Kr], the authors express generating functions for the number of permutations avoiding certain collections T in terms of Chebyshev polynomials. More specifically, the main result of [CW] can be formulated as follows.

Theorem 1.1 ([CW, Theorem 3.1]). *Let $T_1 = \{(1, 2, 3), (k-1, k-2, \dots, 1, k)\}$, $T_2 = \{(2, 1, 3), (1, 2, \dots, k)\}$, and $T_3 = \{(2, 1, 3), (k, 1, 2, \dots, k-1)\}$, then the sequences $|\mathfrak{S}_n(T_i)|$, $i = 1, 2, 3$, have the same generating function*

$$R_k(x) = \frac{U_{k-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_k\left(\frac{1}{2\sqrt{x}}\right)},$$

where $U_r(\cos \theta) = \sin(r+1)\theta/\sin \theta$ is the Chebyshev polynomial of the second kind.

Paper [MV] gives a different proof of Theorem 1.1 for the case of T_2 , and provides explicit expressions for the generating functions of permutations avoiding $(2, 1, 3)$

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

and containing $(1, 2, \dots, k)$ a fixed number of times, or avoiding $(1, 2, \dots, k)$ and containing $(2, 1, 3)$, via the same Chebyshev polynomials. Still another proof of Theorem 1.1 (for all the three cases) based on the use of Dyck paths is given in [Kr], together with the corresponding generalizations (to permutations avoiding the shorter pattern and containing the longer one a fixed number of times).

In this paper, we give a completely different generalization of Theorem 1.1 for the case of T_1 . Following [Bo], we say that $\tau \in \mathfrak{S}_k$ is *p-layered* if it consists of the disjoint union of p substrings (the *layers*) so that the entries decrease within each layer, and increase between the layers. For any $k_1, \dots, k_p \geq 1$ we denote by $w(k_1, \dots, k_p)$ the unique layered pattern of length $k = k_1 + \dots + k_p$ whose i th layer is of length k_i . For example, the only 1-layered pattern of length k is $w(k) = (k, k-1, \dots, 1)$, and the only k -layered pattern of length k is $w(1, \dots, 1) = (1, 2, \dots, k)$. Thus, the longer pattern in the collection T_1 above is just the 2-layered pattern $w(k-1, 1)$.

The main result of this paper is as follows.

Theorem 1.2. *Let $T = \{(1, 2, 3), \tau\}$, where $\tau \in \mathfrak{S}_k$, $k \geq p$, is an arbitrary p -layered pattern, and let $F_T(x) = \sum_{n \geq 0} |\mathfrak{S}_n(T)| x^n$. Then:*

- (i) *if $p = 1$ then $F_T(x)$ is a polynomial of degree $2k - 2$;*
- (ii) *if $p = 2$ then $F_T(x) = R_k(x)$;*
- (iii) *if $p \geq 3$ then $F_T(x) = (1 - \sqrt{1 - 4x})/2x$.*

In fact, only statement (ii) is new, while (i) and (iii) are added for the sake of completeness. Indeed, if $p = 1$ then $\tau = w(k)$, and by [ES], there exists a permutation in $\mathfrak{S}_n$ avoiding both $(1, 2, \dots, l)$ and $(m, m-1, \dots, 1)$ for $n = (m-1)(l-1)$, and there are no such permutations for any $n > (m-1)(l-1)$. On the other hand, if $p \geq 3$ then τ evidently contains $(1, 2, 3)$, hence $|\mathfrak{S}_n(T)| = |\mathfrak{S}_n(1, 2, 3)| = c_n$, the n th Catalan number (see e.g. [Kn, SS]), and the generating function of Catalan numbers is known to be $(1 - \sqrt{1 - 4x})/2x$.

Our proof of Theorem 1.2 is based on finding a recursion for the numbers in question by purely analytical means. In spite of the paradigm formulated in [Kr], that any enumeration problem leading to Chebyshev polynomials is related to Dyck paths, it would be tempting to find a proof that exploits such a relation.

The final version of this paper was written during the second author's (A.V.) stay at Max-Planck-Institut für Mathematik in Bonn, Germany. A.V. wants to express his gratitude to MPIM for the support.

2. PROOFS

Denote by $T^{d,k}$ the collection $\{(1, 2, 3), w(k-d, d)\}$, and by $f_n^{d,k}$ the number of permutations in $\mathfrak{S}_n(T^{d,k})$, $1 \leq d \leq k-1$. The following observation is obtained straightforwardly from the definitions and usual symmetry operations (reversal and complement).

Lemma 2.1. *$f_n^{d,k} = f_n^{k-d,k}$ for all d , $1 \leq d \leq k-1$.*

In what follows we assume that $k \geq 3$ and d are fixed, and omit the indices d, k whenever appropriate; for example, instead of $f_n^{d,k}$ we write just f_n . Moreover, it follows from Lemma 2.1 that we may assume that

$$k - d \geq d. \tag{1}$$

For any $n \geq 1$ and any m such that $1 \leq m \leq n$, we denote by $g_n(i_1, \dots, i_m) = g_n^{d,k}(i_1, \dots, i_m)$ the number of permutations $\pi \in \mathfrak{S}_n(T^{d,k})$ such that $\pi_j = i_j$ for $j = 1, \dots, m$; the corresponding subset of $\mathfrak{S}_n(T^{d,k})$ is denoted by $G_n(i_1, \dots, i_m)$.

The following properties of the numbers $g_n(i_1, \dots, i_m)$ can be deduced easily from the definitions.

Lemma 2.2. (i) *Let $n \geq 2$ and $1 \leq i \leq n$, then*

$$g_n(\dots, i, \dots, i, \dots) = 0.$$

(ii) *Let $n \geq 3$, $1 \leq m \leq n - 2$, and $n - 2 \geq i_1 > i_2 > \dots > i_m \geq 1$, then*

$$g_n(i_1, \dots, i_m, j) = 0 \quad \text{for} \quad i_m + 1 \leq j \leq n - 1.$$

(iii) *Let $n \geq 3$, $1 \leq m \leq \min\{n, k - d\} - 1$, and $n - 1 \geq i_1 > i_2 > \dots > i_m \geq 1$, then*

$$g_n(i_1, \dots, i_m, n) = g_{n-1}(i_1, \dots, i_m).$$

(iv) *Let $n \geq 3$, $2 \leq m \leq n$, $i_1 \geq n - d + 1$, and $n \geq i_1 > i_2 > \dots > i_m \geq 1$, then*

$$g_n(i_1, \dots, i_m) = g_{n-1}(i_2, \dots, i_m).$$

(v) *Let $n \geq k$, $k - d \leq m \leq n - d$, and $n - d \geq i_1 > i_2 > \dots > i_m \geq 1$, then*

$$g_n(i_1, \dots, i_m) = 0.$$

Proof. Property (i) is evident. To prove (ii) is enough to observe that if $\pi \in G_n(i_1, \dots, i_m, j)$ then the entries i_m, j , and n give an occurrence of $(1, 2, 3)$ in π , a contradiction.

To prove (iii), denote by π^* the permutation obtained from π by deleting its largest entry. Then $\pi \in G_n(i_1, \dots, i_m, n)$ if and only if $\pi^* \in G_{n-1}(i_1, \dots, i_m)$, since entry n placed as in (iii) cannot be used in an occurrence of $(1, 2, 3)$ or $w(k - d, d)$ in π .

The proof of (iv) goes along the same lines. We denote by π° the permutation obtained from π by deleting entry i_1 and subtracting 1 from all the entries greater than i_1 ; then $\pi \in G_n(i_1, \dots, i_m)$ if and only if $\pi^\circ \in G_{n-1}(i_2, \dots, i_m)$. Indeed, the only if part is trivial, so assume that $\pi^\circ \in G_{n-1}(i_2, \dots, i_m)$, and there exists an occurrence of $(1, 2, 3)$ or $w(k - d, d)$ in π . Evidently, such an occurrence makes use of i_1 . If (i_1, j_1, j_2) is an occurrence of $(1, 2, 3)$, then j_1 and j_2 lie to the right of i_m , and since $i_1 > i_m$, the triple (i_m, j_1, j_2) is an occurrence of $(1, 2, 3)$ in π° , a contradiction. Otherwise, i_1 enters an occurrence of $w(k - d, d)$, and hence π contains at least d entries strictly greater than i_1 , a contradiction.

Finally, to prove (v), consider all the entries of π greater than i_1 . There are at least d such entries, and all of them lie to the right of i_m . If a pair (j_1, j_2) of these entries is an occurrence of $(1, 2)$, then (i_m, j_1, j_2) is an occurrence of $(1, 2, 3)$, a contradiction. Hence, all these entries provide an occurrence of $(p, p - 1, \dots, 1)$ for some $p \geq d$; therefore, any d out of these entries, together with any $k - d$ entries out of $i_1, \dots, i_m$, give an occurrence of $w(k - d, d)$. $\square$

Now we introduce several objects that play an important role in the proof of the main result. For $n \geq d + 1$ and $1 \leq m \leq \min\{n, k\} - d$ we define

$$\begin{aligned}\mathcal{I}(n, m) &= \{(i_1, \dots, i_m) : m - j \leq i_{j+1} \leq i_j - 1, j = 0, \dots, m - 1\}, i_0 = n - d + 1, \\ \mathcal{J}(n, m) &= \{(i_1, \dots, i_m) : m - j \leq i_{j+1} \leq i_j - 1, j = 0, \dots, m - 1\}, i_0 = n - d + 2, \\ \mathcal{K}(n, m) &= \{(i_1, \dots, i_m) : (i_1, \dots, i_{m-1}) \in \mathcal{I}(n, m - 1), 1 \leq i_m \leq i_{m-1} - 1\},\end{aligned}$$

and put

$$A(n, m) = \sum_{(i_1, \dots, i_m) \in \mathcal{I}(n, m)} g_n(i_1, \dots, i_m).$$

Theorem 2.3. *Let $n \geq d + 2$ and $2 \leq m \leq \min\{n, k - d\} - 1$, then*

$$A(n, m) = A(n - 1, m - 1) + A(n, m + 1).$$

Proof. By the definition,

$$A(n, m) = \sum_{(i_1, \dots, i_m) \in \mathcal{I}(n, m)} \sum_{j=1}^n g_n(i_1, \dots, i_m, j).$$

Let us decompose the internal sum into three subsums over the intervals $1 \leq j \leq i_m - 1$, $i_m \leq j \leq n - 1$, $j = n$, and denote them S_1 , S_2 , and S_3 , respectively. Evidently,

$$\begin{aligned}S_1 &= \sum_{(i_1, \dots, i_m) \in \mathcal{I}(n, m)} \sum_{j=1}^{i_m-1} g_n(i_1, \dots, i_m, j) \\ &= \sum_{(i_1, \dots, i_m, j) \in \mathcal{K}(n, m+1)} g_n(i_1, \dots, i_m, j).\end{aligned}$$

It is easy to see that if $(i_1, \dots, i_m, j) \in \mathcal{K}(n, m + 1)$ and $(i_1, \dots, i_m, j) \notin \mathcal{I}(n, m + 1)$ then there exists p , $1 \leq p \leq m$, such that $i_p = j$, and therefore by Lemma 2.2(i),

$$S_1 = \sum_{(i_1, \dots, i_m, j) \in \mathcal{I}(n, m+1)} g_n(i_1, \dots, i_m, j) = A(n, m + 1).$$

Next,

$$S_2 = \sum_{(i_1, \dots, i_m) \in \mathcal{I}(n, m)} \sum_{j=i_m}^{n-1} g_n(i_1, \dots, i_m, j) = 0$$

by Lemma 2.2(i) and (ii). Finally, by Lemma 2.2(iii),

$$\begin{aligned}S_3 &= \sum_{(i_1, \dots, i_m) \in \mathcal{I}(n, m)} g_n(i_1, \dots, i_m, n) \\ &= \sum_{(i_1, \dots, i_m) \in \mathcal{J}(n-1, m)} g_{n-1}(i_1, \dots, i_m).\end{aligned}$$

Denote the latter expression by $B(n-1, m)$, so

$$A(n, m) = A(n, m+1) + B(n-1, m), \quad (2)$$

for $n \geq d+1$, $1 \leq m \leq \min\{n, k-d\} - 1$.

Consider now $B(n, m+1)$. One has

$$\begin{aligned} B(n, m+1) &= \sum_{(i_1, \dots, i_m, j) \in \mathcal{I}(n, m+1)} g_n(i_1, \dots, i_m, j) \\ &+ \sum_{(i_2-1, \dots, i_m-1, j-1) \in \mathcal{J}(n, m)} g_n(n-d+1, i_2, \dots, i_m, j), \end{aligned}$$

and by Lemma 2.2(iv), the second term in the right hand side equals $B(n-1, m)$. Therefore,

$$B(n, m+1) = A(n, m+1) + B(n-1, m)$$

for $n \geq d+1$, $1 \leq m \leq \min\{n, k-d\} - 1$.

Comparing this with (2) one gets $B(n, m+1) = A(n, m)$ for $n \geq d+1$, $1 \leq m \leq \min\{n, k-d\} - 1$, and hence

$$A(n, m) = A(n-1, m-1) + A(n, m+1)$$

for $n \geq d+2$, $2 \leq m \leq \min\{n, k-d\} - 1$. $\square$

Our next aim is to express $A(n, m)$ in terms of f_j 's. We start from the following result.

Lemma 2.4. *Let $n \geq d+2$ and $1 \leq j \leq d$, then*

$$g_n(n+1-j) = \sum_{i=0}^{j-1} (-1)^i \binom{j-1-i}{i} f_{n-1-i}. \quad (3)$$

Proof. Let $n \geq 3$; it is clear that $g_n(n) = f_{n-1}$, hence (3) holds for $j=1$ and $n \geq 3$. Moreover, if $d \neq 1$ then $g_n(n-1) = f_{n-1}$ as well, and hence (3) holds also for $j=2$ and $n \geq 3$. This gives the basis for the induction.

Next, observe that

$$g_n(n+2-j) - g_n(n+1-j) = g_{n-1}(n+2-j)$$

for $n \geq 3$, $3 \leq j \leq \min\{d, n\}$. Indeed,

$$g_n(r) = \sum_{i=1}^n g_n(r, i).$$

The sum in the right hand side is subdivided into four sums: for $1 \leq i \leq r-1$, for $i=r$, for $r+1 \leq i \leq n-1$, and for $i=n$, respectively. Using Lemma 2.2(iv), (i), (ii), and (iii) respectively, we get

$$g_n(r) = \sum_{i=1}^r g_{n-1}(i)$$

for $n \geq 3$ and $n + 1 - d \leq r \leq n - 1$. Observe that condition $m \leq k - d - 1$ from Lemma 2.2(iii) is satisfied by virtue of (1) and $k \geq 3$. It remains to substitute $r = n - j + 1$ and $r = n - j + 2$, and to find the difference between the right hand sides; the restrictions on r are transformed into $3 \leq j \leq d$.

Finally, it is easy to check that

$$\begin{aligned} \sum_{i=0}^{j-2} (-1)^i \binom{j-2-i}{i} f_{n-1-i} - \sum_{i=0}^{j-3} (-1)^i \binom{j-3-i}{i} f_{n-2-i} \\ = \sum_{i=0}^{j-1} (-1)^i \binom{j-1-i}{i} f_{n-1-i}, \end{aligned}$$

for any $n \geq 2$ and $j \geq 3$, and hence (3) is satisfied for $n \geq 3$ and $1 \leq j \leq \min\{d, n-1\}$. It follows now from (1) that (3) is satisfied for $n \geq d+2$ and $1 \leq j \leq d$. $\square$

We are now ready to find an expression for $A(n, m)$ similar to expression (3) for $g_n(n+1-j)$.

Theorem 2.5. *Let $n \geq k$ and $1 \leq m \leq k - d$, then*

$$A(n, m) = \sum_{i=0}^{m+d} (-1)^i \binom{m+d-i}{i} f_{n-i}. \quad (4)$$

Proof. Similarly to the proof of Lemma 2.4, we first check (4) for $m = 1$ and $m = 2$. Clearly,

$$A(n, 1) = \sum_{j=1}^{n-d} g_n(j) = f_n - \sum_{j=1}^d g_n(n+1-j).$$

Using (3) we get

$$\begin{aligned} A(n, 1) &= f_n - \sum_{j=1}^d \sum_{i=0}^{j-1} (-1)^i \binom{j-1-i}{i} f_{n-1-i} \\ &= f_n - \sum_{i=0}^{d-1} (-1)^i f_{n-1-i} \sum_{j=i+1}^d \binom{j-1-i}{i} \\ &= \sum_{i=0}^{d+1} (-1)^i \binom{d+1-i}{i} f_{n-i} \end{aligned}$$

for $n \geq d+2$, so (4) holds for $m = 1$ and $n \geq d+2$.

Next, by (2), we have

$$A(n, 2) = A(n, 1) - B(n-1, 1) = A(n, 1) - A(n-1, 1) - g_{n-1}(n-d)$$

for $n \geq d + 2$ (condition $m = 1 \leq k - d - 1$ follows from (1) and $k \geq 3$ as in the proof of Lemma 2.4). Using (3) for $j = d$ and (4) for $m = 1$ we get

$$\begin{aligned} A(n, 2) &= \sum_{i=0}^{d+1} (-1)^i \binom{d+1-i}{i} f_{n-i} \\ &\quad - \sum_{i=0}^{d+1} (-1)^i \binom{d+1-i}{i} f_{n-1-i} - \sum_{i=0}^{d-1} (-1)^i \binom{d-1-i}{i} f_{n-2-i} \\ &= \sum_{i=0}^{d+2} (-1)^i \binom{d+2-i}{i} f_{n-i} \end{aligned}$$

for $n \geq d + 2$, so (4) holds for $m = 2$ and $n \geq d + 2$.

Finally, by Theorem 2.3,

$$A(n, m) = A(n, m-1) - A(n-1, m-2)$$

for $n \geq d + 2$ and $3 \leq m \leq k - d$, so by induction we get

$$\begin{aligned} A(n, m) &= \sum_{i=0}^{m-1+d} (-1)^i \binom{m-1+d-i}{i} f_{n-i} - \sum_{i=0}^{m-2+d} (-1)^i \binom{m-2+d-i}{i} f_{n-1-i} \\ &= \sum_{i=0}^{m+d} (-1)^i \binom{m+d-i}{i} f_{n-i} \end{aligned}$$

for $n \geq d + 2$ and $1 \leq m \leq \min\{n+1, k\} - d$, and hence for $n \geq k$ and $1 \leq m \leq k - d$. $\square$

Now we are ready to prove Theorem 1.2 for $k \geq 3$. First of all, it follows from Lemma 2.2(v) that $A(n, k-d) = 0$ for all $n \geq k$. Hence, by Theorem 2.5,

$$\sum_{i=0}^k (-1)^i \binom{k-i}{i} f_{n-i} = 0$$

for $n \geq k$, or, equivalently,

$$\sum_{i=0}^k \sum_{n \geq k} (-1)^i x^i \binom{k-i}{i} f_{n-i} x^{n-i} = 0.$$

Evidently, for $n \leq k - 1$ the layered restriction is void, so by [Kn, SS], $f_n = c_n$ for $0 \leq n \leq k - 1$, where c_n is the n th Catalan number. Therefore

$$\sum_{i=0}^k (-x)^i \binom{k-i}{i} \left(F_T(x) - \sum_{j=0}^{k-i-1} x^j c_j \right) = 0. \quad (5)$$

Recall that the Chebyshev polynomials of the second kind satisfy relation

$$x^{k/2} U_k \left(\frac{1}{2\sqrt{x}} \right) = \sum_{i=0}^k (-x)^i \binom{k-i}{i}$$

(see [Ri, pp. 75,76]), while the Catalan numbers satisfy relation

$$\sum_{i=0}^l (-1)^i \binom{k-i}{i} c_{l-i} = (-1)^l \binom{k-1-l}{l}$$

for $l \leq k-1$ (see [Ri, pp. 152–154]), and hence

$$\sum_{i=0}^k (-x)^i \binom{k-i}{i} \sum_{j=0}^{k-i-1} x^j c_j = \sum_{l=0}^{k-1} x^l \sum_{i=0}^l (-1)^i \binom{k-i}{i} c_{l-i} = \sum_{l=0}^{k-1} (-x)^l \binom{k-1-l}{l}.$$

Therefore, (5) yields

$$F_T(x) = \frac{\sum_{i=0}^{k-1} (-x)^i \binom{k-1-i}{i}}{\sum_{i=0}^k (-x)^i \binom{k-i}{i}} = \frac{U_{k-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_k\left(\frac{1}{2\sqrt{x}}\right)} = R_k(x)$$

for $k \geq 3$.

The remaining case $k = 2$ is trivial, since $R_2(x) = (1-x)^{-1}$, which is equivalent to the fact that there is exactly one permutation of length n avoiding $w(1, 1) = (1, 2)$.

REFERENCES

- [Bo] M. Bóna, *The solution of a conjecture of Stanley and Wilf for all layered patterns*, J. Combin. Theory Ser. A **85** (1999), 96–104.
- [CW] T. Chow and J. West, *Forbidden subsequences and Chebyshev polynomials*, Discr. Math. **204** (1999), 119–128.
- [ES] P. Erdős and G. Szekeres, *A combinatorial problem in geometry*, Compositio Mathematica **2** (1935), 463–470.
- [Kn] D. Knuth, *The Art of Computer Programming*, vol. 3, Addison Wesley, Reading, MA, 1973.
- [Kr] C. Krattenthaler, *Permutations with restricted patterns and Dyck paths* (2000), preprint CO/0002200.
- [MV] T. Mansour and A. Vainshtein, *Restricted permutations, continued fractions, and Chebyshev polynomials*, Electron. J. Combin. **7** (2000), #R17.
- [Ri] J. Riordan, *Combinatorial Identities*, John Wiley, New York, 1968.
- [SS] R. Simion and F. Schmidt, *Restricted permutations*, European J. Combin. **6** (1985), 383–406.